

Home Remedies

Animal General of Cranberry

find on our facebook page under **Notes**

Fear of Thunder Storms

ACONITE (Homeopathic aconite)

Use: Aconite is a pill used in dogs to calm their fears of storms, fireworks, etc. It can be purchased on-line or at GNC stores. It comes in 6C, 12C or 30C - all three sizes are safe to use and are non-toxic.

Dose: Small dogs (1-15 lb) Give 4 tablets twice daily

Medium dogs (16-50 lb) Give 6 tablets twice daily

Large dogs (over 50 lbs) Give 8 tablets twice daily

Family Member is Allergic to Cat or Dog?

ALLERGY RINSE

Use: This is a rinse that is applied to cats or dogs when the pet owner is allergic to that pet.

Method : First, mix 9 parts warm water with 1 part white vinegar into a gallon pail. Sponge over the dog or cat thoroughly. Then, mix 1 gallon of warm water and 1 capful of Downey fabric softener. Sponge mixture

over the pet, do NOT rinse. Let air dry. This can be applied every 2 to 6 wks.

ALLERGY TREATMENT FOR DOGS

Benadryl

Use: This is an over-the-counter anti-histamine that can be used to decrease mild allergy signs in dogs, such as itching and congestion. It comes in a liquid children's elixer (12.5 mg /tsp) or in 25 mg or 50 mg capsules or pills.

Dose: Use 1 mg/lb by mouth up to 3 times/day as needed.

ALLERGY TREATMENT FOR CATS

Chlorpheniramine

Use: This is an over-the-counter anti-histamine that can be used to decrease mild allergy signs in cats, such as itching and congestion. It comes in 4 mg tablets.

Dose: Give 1/2 of a 4 mg tablet by mouth up to twice daily as needed.

BEE STINGS/INSECT BITES - Canine

Benadryl

Use: This is an over-the-counter anti-histamine that can be used for hives, facial swelling and inflammation that may occur after a dog has been stung by a bee or spider.

Dose: Use 1 mg/lb by mouth up to 3 times/day as needed.

Ice pack wrapped in a small towel can also be applied to the affected skin. Call your vet

immediately and tell them you gave Benadryl.

Canine

BLAND DIET

Canine bland diet:

Use: A bland diet can be offered to a dog that has mild gastro-intestinal upset. This diet is meant to be fed for a few days only.

Method: Mix 3 parts boiled rice with 1 part cooked chicken or lean ground beef. It is best to feed small, frequent meals to a dog that has an upset stomach. Gradually reintroduce your pet to his regular diet by mixing the bland diet with the regular diet over a few days.

Feline

bland diet:

Use: A bland diet can be offered to a cat that has mild gastro-intestinal upset. This diet is meant to be fed for a few days only.

Method: Plain meat baby food (such as Gerber's chicken or turkey) fed 1-2 teaspoons at a time, warmed.

EAR CLEANER To Prevent Infections

Mix 2 parts rubbing alcohol with 1 part white vinegar. This mixture can be instilled into a dog's ear.

Gently massage the ear solution into the ear for 15-30 seconds, then wipe out excess with a cotton ball. This can be

repeated no more than every other day as needed. Do NOT use this solution if the ear is inflamed or sore - the rubbing alcohol will sting!

NAIL BLEEDING TREATMENT

If you cut a nail too short, it will bleed. Prior to cutting your pets nails, it's a good idea to have Quik-stop (available at pet stores) on hand. Apply Quik-stop directly to the bleeding nail with direct pressure until the bleeding stops. If you don't have Quik-stop, corn starch will also work as well as a tea bag.

SKUNK DE-ODORIZER

Mix 1 quart of 3% hydrogen peroxide + 1/4 cup of baking soda + 1 teaspoon of liquid dish soap (Dawn dish soap works well) -mix the ingredients and use immediately. The chemical reaction produced from these ingredients lasts only a limited time.

Do NOT wet the dog first!!!

Pour the solution over your pet, rub it in thoroughly (remember to wear gloves so you don't smell

like the skunk!). Then let the pet soak for a few minutes. RINSE off. Do not leave the solution on your pet. Repeat if necessary

TOXIN INGESTION TREATMENT

If you have witnessed your dog eating something toxic (medicine, chocolate, onions), you may induce vomiting by giving Hydrogen peroxide.

Dose: 1 tablespoon / 15 lb of body weight. Repeat in 15 minutes if the dog hasn't vomited.

Use: Induce vomiting within 30-60 minutes of ingestion of the toxic substance. If it has been longer than 60 minutes, it will not work.

Warning: If the pet has ingested anything caustic/ flammable - do NOT induce vomiting

Pittsburgh Poison Control : 412-681-6669